

# The Weight Performance Stability of Mice on Modeling Obesity-Associated Hyperglycemia Induced by Dextrose Monohydrate

Deksa Yudha Syach Putra<sup>1</sup>, Setiyo Budi Santoso<sup>1,2,\*</sup>, Heni Lutfiyati<sup>1,2</sup>

<sup>1</sup>Department of Pharmacy, Faculty of Health Sciences; <sup>2</sup>Consortium for Pharmacology and Clinical Pharmacy Studies, Universitas Muhammadiyah Magelang, Indonesia.

Corresponding author\*

sb@unimma.ac.id

Manuscript received: 11 August, 2022. Revision accepted: 26 August, 2022. Published: 06 September, 2022.

## Abstract

Previously, streptozotocin and alloxan were employed to imitate hyperglycemia in mice. High doses of sucrose were also induced as an alternative. Due to body mass index has been associated with hyperglycemia, the evidence of weight body index in various induction alternate kinds, however, have not been fully reported. Here-in, we report on the weight performance stability of mice body weight induced by dextrose, streptozotocin, and alloxan. To begin, all mice were divided into six groups of five, with one reserve in each. Following seven days of acclimatization, the mice were induced for nine days of hyperglycemia modeling; alloxan (Groups A and D), streptozotocin (Group B and E), dextrose monohydrate (Groups C and F). On preclinical research animals modelling related to obesity-associated hyperglycemia in mice, dextrose monohydrate induction was most successful than streptozotocin and alloxan induction, which performed best during the induction period (31% weight growth) and after metformin intervention (36% weight growth). Overall, dextrose monohydrate is most suitable to be used for modeling type 2 diabetes mellitus test animals rather than alloxan and streptozotocin.

**Keywords:** alloxan; preclinical research animals; streptozotocin; type 2 diabetes mellitus.

**Abbreviations:** type 1 diabetes mellitus (T1DM), type 2 diabetes mellitus (T2DM), glucose transporter 2 (GLUT2), body weight (BW)

## INTRODUCTION

Diabetic mice are commonly employed in the development of anti-diabetes agent. Alloxan and streptozotocin are commonly used to induce diabetic in the mice (Al-awar et al., 2016; King & Austin, 2017; Kottaisamy et al., 2021). In a single induction, alloxan elevated the glycemic index to 127 mg/dL (Irdalisa et al., 2015), whereas three days or more raised it to 156 - 270 mg/dL (Dewi et al., 2021; Hamdani & Nurman, 2020; Lolok et al., 2019). Streptozotocin had a comparable impact, increasing glucose levels to 136 mg/dL after three days (Apriani et al., 2011) and 220-244 mg/dL after five days (Ocktarini et al., 2011; Suwanto & Rahmawati, 2019).

As an alternative to alloxan and streptozotocin, glucose induction groups have been induced. Fructose intake for 12 weeks hold glycemic levels at 81 mg/dL (Tillman et al., 2014). Dextrose outperformed fructose after seven days of induction, while by 10%-concentration raised glycemic at 148 mg/dL

(Pramushinta et al., 2019) and by 40% at 154 mg/dL (Santoso & Suryanto, 2017).

Diabesity is a new term for diabetes that comes alongside obesity (Serván, 2013). Obese are roughly three times more vulnerable to diabetic (Prasetyani & Sodikin, 2017). Diabetes, most especially type 2 diabetes mellitus (T2DM), is correlated with weight gained (Droz et al., 2017) or obesity (Fang et al., 2019; Finkelstein et al., 2012). Weight gained and body mass are two factors that contribute to rising glycemic levels (Algoblan et al., 2014), so obesity and T2DM called the twin pandemic (Scheithauer et al., 2016).

Previously, streptozotocin and alloxan were employed to imitate hyperglycemia in mice. High doses of sucrose were also induced as an alternative. Due to body mass index has been associated with hyperglycemia, the evidence of weight body index in various induction alternate kinds, however, have not been fully reported. Here-in, we report on the weight performance stability of mice body weight caused by dextrose, streptozotocin, and alloxan.

## MATERIALS AND METHODS

### Ethical Clearance

The Medical and Health Research Ethics Committee Ethics of the Faculty of Medicine, Universitas Gadjah Mada, has accepted the whole series of research procedures under registration number KE/FK/0328/EC/2022.

### Animals

Male Balb/c mice were around 2-3 months that bred by Pharmacy Laboratory of Universitas Gadjah Mada (Yogyakarta, Indonesia) and treated in accordance with current norms.

### Materials

Alloxan monohydrate (Sigma-Aldrich), streptozotocin (Sigma-Aldrich), and dextrose monohydrate 40% (Otsuka) were used to induce mice. Metformin (Dexa Medica) and Natrium Carboxy Methylcellulose (Dai-ichi Kogyo Seiyaku) were prepared to assess the stability of hyperglycemia treatment. Citrate buffer pH 4.5 is solved for intraperitoneal preparation. Supplies for glycemic testing, such as strip and glucometer (easytouch). Among the other consumables are a handsoon, syringe for 1mL and 5 mL, blood lancet, and aquadest. As support equipment, analytical scales, beakers, mouse cages, markers, mortars and stampers, and oral probes were used.

### Procedures

All mice were divided into six groups of five, with one reserve in each. Following seven days of acclimatization, the mice were induced for nine days of hyperglycemia modeling and five days of metformin intervention.

#### ■ Induction Period

To get baseline data, we assessed the weight of the mice following the acclimatization. For 9 days, Groups A and D were induced alloxan 0.12 mg/gram body weight (BW) intraperitoneally, Group B and E were induced streptozotocin 0.05 mg/gram BW intraperitoneally, and Groups C and F were induced dextrose monohydrate 6 mg/gram BW orally. At the end of the induction, we weighed the mice again.


#### ■ Intervention Period

Following the induction period, groups A, B, and C were treated metformin orally, whereas groups D, E, and F as negative controls and were treated Na CMC orally. Dextrose monohydrate continued to activate Groups C and F throughout the intervention. On day 14, the body weight was measured.

## RESULTS AND DISCUSSION

### Performance of Alloxan Induction


Mice in groups A and D gained 35.86 grams and 37.98 grams of body weight after nine days of alloxan induction, respectively. Mice on Group A grew to 37.50 grams after a five-day metformin intervention, whereas control mice (Group D) stayed constant at 37.98 grams. (Figure 1).



**Figure 1.** The body weight gain due to alloxan induction in the intervention group (A) and the control group (D). The solid line represents the induction period, the dashed line represents the intervention period.

### Performance of Streptozotocin Induction

Mice in groups B and E gained 35.80 grams and 36.60 grams of body weight after nine days of streptozotocin induction, respectively. Mice on Group B grew to 39.10 grams after a five-day metformin intervention, whereas control mice (Group D) stayed constant at 36.60 grams. (Figure 2).




**Figure 2.** The body weight gain due to streptozotocin induction in the intervention group (B) and the control group (E). The solid line represents the induction period, the dashed line represents the intervention period.

### Performance of Dextrose Monohydrate Induction

Mice in groups C and F gained 40.33 grams and 36.08 grams of body weight after nine days of dextrose monohydrate induction, respectively. Both groups thereafter continued to rise to 41.90 grams and 39.36

grams respectively, after receiving metformin intervention for five days (Figure 3).




**Figure 3.** The body weight gain due to dextrose monohydrate induction in the intervention group (C) and the control group (F). The solid line represents the induction period, the dashed line represents the intervention period.

**Induction variant performance in mice treated with metformin was compared.**

After the 9-day induction period, dextrose-induced mice gained the most weight (40.33 grams), followed by alloxan-induced mice (35.86 grams), and streptozotocin-induced mice (35.80 grams). After 5 days of metformin treatment, dextrose-induced mice had the best body weight performance (41.90 grams), followed by

streptozotocin-induced (39.10 grams) and alloxan-induced (37,50 gram)



**Figure 4.** The body weight gain in mice following 9 days of induction (solid line) to alloxan (A), streptozotocin (B), and dextrose monohydrate (C), and persistence rate following 5 days of metformin intervention (dash line) after obesity (day 14<sup>th</sup>).

Dextrose monohydrate induction in mice was most successful than streptozotocin and alloxan induction, which performed best during the induction period (31% weight growth) and after metformin intervention (36% weight growth).

**Table 1.** Percentage of body weight gain in mice following 9 days of induction to alloxan (A), streptozotocin (B), and dextrose monohydrate (C), and persistence rate following 5 days of metformin intervention after obesity (day 14<sup>th</sup>).

Group Label	Induction Treatment	Body Weight (Percentage Increase)		
		Baseline	9th day treatment	14th day treatment
A	Alloxan	31.34±1.27 (-)	35.86±3.58 (14%)	37.50±4.54 (20%)
B	Streptozotocin	31.86±2.69 (-)	35.80±3.58 (12%)	39.10±3.01 (23%)
C	Dekstrose Monohydrate	30.88±2.35 (-)	40.33±2.66 (31%)	41.90±3.28 (36%)

**Discussion**

On preclinical research animals, we simulate obesity-associated hyperglycemia in mice using dextrose monohydrate induction. The bulk previous studies employed streptozotocin and alloxan to imitate hyperglycemia. However, as one of the important criteria in hyperglycemia etiology, the investigation of body weight issues was disregarded. Due to the importance of publishing our findings, we would show the evidence of steady rates of dextrose monohydrate-induced body weight during the experiment.

According to the availability of test animals in preclinical research on the developing of anti-diabetic agents, there are two options: genetic modification, or chemical induction (Kottaisamy et al., 2021). In the unavailability of genetically diabetic test animals, alloxan and streptozotocin are widely used to cause

diabetes in mice (Al-awar et al., 2016; Kottaisamy et al., 2021).

Streptozotocin and alloxan, on the other hand, act on beta cells via the glucose transporter 2 (GLUT2) and cause complete ablation of beta cells inside the islets, leading in severe insulin insufficiency, hyperglycemia, and weight loss, they are more suited for modeling Type 1 diabetes (King & Austin, 2017). Mice given alloxan or streptozotocin lost weight because they were unable to use the available glucose for energy. To meet the body's energy requirements, excessive catabolism of protein and fat in muscle and adipose tissue was performed (Malik et al., 2015; Sinata & Arifin, 2016; Suwanto & Rahmawati, 2019).

Since the great majority of T2DM are overweight (Finkelstein et al., 2012; Wilding, 2014), body mass index has long been linked to hyperglycemia and T2DM

(Algoblan et al., 2014; Jung & Choi, 2014), with the potential of obesity to promote insulin resistance being the primary explanation (Jung & Choi, 2014). Obesity was proclaimed in mice as test animals when they weighed more than 20% of their age-matched mates. (Patonah et al., 2018).

Common sugar exacerbates obesity and increases adiposity in mice (Kleinert et al., 2018). Adipocyte hypertrophy and hyperplasia lead to excessive lipid accumulation (lipotoxicity), which activate insulin resistance (Lee et al., 2020; Longo et al., 2019). Indeed, the diets manipulated of mice with high sucrose feeding generate a compensatory response from some beta cells, (King & Austin, 2017). Additionally, the weight gain was also linked to the dysfunction of leptin (anorexigenic hormone) due to the circulating of glucose which led the test animals to consume more feed (Tillman et al., 2014).

Throughout the investigation, the induction performance of dextrose monohydrate has been shown to increase glycemia. Dextrose monohydrate is more suitable to be used for modeling type 2 diabetes mellitus test animals due to the aspect of increasing body weight, rather than the two toxic compounds that actually suppress body weight, and they are more suitable for modeling mice with diabetes mellitus type 1 as well, aside from having a higher level of glucose elevation than alloxan and streptozotocin.

## CONCLUSIONS

On preclinical research animals modelling related to obesity-associated hyperglycemia in mice, dextrose monohydrate induction was most successful than streptozotocin and alloxan induction, which performed best during the induction period (31% weight growth) and after metformin intervention (36% weight growth). Throughout the investigation, dextrose monohydrate is most suitable to be used for modeling type 2 diabetes mellitus test animals rather than alloxan and streptozotocin.

**Acknowledgements:** This study is part of the acceleration research scheme for institutional vision revitalization research output (reference number: 012/Kontrak/PRVI-PA/2022) in the subject of exploring the biological and pharmacological effects of employing plants, fungi, animals, microbes, and minerals.

**Authors' Contributions:** Conceptualization, SBS; Methodology, SBS and HL; Data Collection, DYSP; Writing-Original Draft Preparation, SBS and DYSP; Writing-Review & Editing, DYSP, SBS, and HL. All authors read and approved the final version of the manuscript.

**Competing Interests:** The authors claim to have no conflicts of interest. The funders had no say in the design of the investigation, data collection, analysis, or interpretation, article writing, or decision to publish the findings. The authors declare that there are no competing interests.

**Funding:** The authors are thankful to the Research, Development, and Community Service Department of Universitas Muhammadiyah Magelang for financial supporting.

## REFERENCES

- Al-awar, A., Kupai, K., Veszelka, M., Szűcs, G., Attieh, Z., Murlasits, Z., Török, S., Pósa, A., & Varga, C. (2016). Experimental Diabetes Mellitus in Different Animal Models. *Journal of Diabetes Research*, 2016, 1–12. <https://doi.org/10.1155/2016/9051426>
- Algoblan, A., Alalfi, M., & Khan, M. (2014). Mechanism linking diabetes mellitus and obesity. *Diabetes, Metabolic Syndrome and Obesity: Targets and Therapy*, 2014(7), 587. <https://doi.org/10.2147/DMSO.S67400>
- Apriani, N., Suhartono, E., & Akbar, I. Z. (2011). Korelasi Kadar Glukosa Darah dengan Kadar Advanced Oxidation Protein Products (AOPP) Tulang pada Tikus Putih Model Hiperqlikemia. *Jurnal Kesehatan Masyarakat*, 11(1), 8.
- Dewi, R. S., Rahayu, L., & Atika, I. (2021). Efek Penurunan Kadar Glukosa Darah Rebusan Asparagus (*Asparagus officinalis* L.) pada Mencit yang Diinduksi Aloksan. *Jurnal Ilmu Kefarmasian Indonesia*, 19(1), 6.
- Droz, B. A., Sneed, B. L., Jackson, C. V., Zimmerman, K. M., Michael, M. D., Emmerson, P. J., Coskun, T., & Peterson, R. G. (2017). Correlation of disease severity with body weight and high fat diet in the FATZO/Pco mouse. *PLOS ONE*, 12(6), e0179808. <https://doi.org/10.1371/journal.pone.0179808>
- Fang, J.-Y., Lin, C.-H., Huang, T.-H., & Chuang, S.-Y. (2019). In Vivo Rodent Models of Type 2 Diabetes and Their Usefulness for Evaluating Flavonoid Bioactivity. *Nutrients*, 11(3), 530. <https://doi.org/10.3390/nu11030530>
- Finkelstein, E. A., Khavjou, O. A., Thompson, H., Trogdon, J. G., Pan, L., Sherry, B., & Dietz, W. (2012). Obesity and Severe Obesity Forecasts Through 2030. *American Journal of Preventive Medicine*, 42(6), 563–570. <https://doi.org/10.1016/j.amepre.2011.10.026>
- Hamdani, I., & Nurman, S. (2020). Ekstrak Etanol Kopi Hijau Arabika (*Coffea arabica* L.) sebagai Antihiperqlikemi pada Mencit (*Mus musculus*). *Jurnal Kefarmasian Indonesia*, 10(2), 140–147. <https://doi.org/10.22435/jki.v10i2.2122>
- Irdalisa, Safrida, Khairil, Abdullah, & Sabri, M. (2015). Profil Kadar Glukosa Darah Pada Tikus Setelah Penyuntikan Aloksan Sebagai Hewan Model Hiperqlikemik. *Jurnal EduBio Tropika*, 3(1).
- Jung, U., & Choi, M.-S. (2014). Obesity and Its Metabolic Complications: The Role of Adipokines and the Relationship between Obesity, Inflammation, Insulin Resistance, Dyslipidemia and Nonalcoholic Fatty Liver Disease. *International Journal of Molecular Sciences*, 15(4), 6184–6223. <https://doi.org/10.3390/ijms15046184>

- King, A., & Austin, A. (2017). Chapter 10 Animal Models of Type 1 and Type 2 Diabetes Mellitus. In *Animal Models for the Study of Human Disease* (pub.1086151127; pp. 245–265). <https://doi.org/10.1016/b978-0-12-809468-6.00010-3>
- Kleinert, M., Clemmensen, C., Hofmann, S. M., Moore, M. C., Renner, S., Woods, S. C., Huypens, P., Beckers, J., de Angelis, M. H., Schürmann, A., Bakhti, M., Klingenspor, M., Heiman, M., Cherrington, A. D., Ristow, M., Lickert, H., Wolf, E., Havel, P. J., Müller, T. D., & Tschöp, M. H. (2018). Animal models of obesity and diabetes mellitus. *Nature Reviews Endocrinology*, *14*(3), 140–162. <https://doi.org/10.1038/nrendo.2017.161>
- Kottaisamy, C. P. D., Raj, D. S., Prasanth Kumar, V., & Sankaran, U. (2021). Experimental animal models for diabetes and its related complications—A review. *Laboratory Animal Research*, *37*(1), 23. <https://doi.org/10.1186/s42826-021-00101-4>
- Lee, S., Goodson, M., Vang, W., Kalanetra, K., Barile, D., & Raybould, H. (2020). 2'-Fucosyllactose Supplementation Improves Gut-Brain Signaling and Diet-Induced Obese Phenotype and Changes the Gut Microbiota in High Fat-Fed Mice. *Nutrients*, *12*(4), 1003. <https://doi.org/10.3390/nu12041003>
- Lolok, N., Rahmat, H., & Wijayanti, P. M. (2019). Efek Antidiabetes Kombinasi Ekstrak Kulit Bawang Dayak Dan Kulit Bawang Merah Pada Mencit Yang Diinduksi Aloksan. *Jurnal Mandala Pharmacon Indonesia*, *5*(2), 9.
- Longo, M., Zatterale, F., Naderi, J., Parrillo, L., Formisano, P., Raciti, G. A., Beguinot, F., & Miele, C. (2019). Adipose Tissue Dysfunction as Determinant of Obesity-Associated Metabolic Complications. *International Journal of Molecular Sciences*, *20*(9), 2358. <https://doi.org/10.3390/ijms20092358>
- Malik, M. I., Nasrul, E., & Asterina, A. (2015). Hubungan Hiperglikemia dengan Prothrombin Time pada Mencit (*Mus musculus*) yang Diinduksi Aloksan. *Jurnal Kesehatan Andalas*, *4*(1). <https://doi.org/10.25077/jka.v4i1.219>
- Octarini, R., Prasetyo, D. H., & Sjarifah, I. (2011). Effect of herbal extract of anting-anting (*Acalypha australis*) on blood glucose level of Balb/C mice with induction of Streptozotocin. *Biofarmasi Journal of Natural Product Biochemistry*, *9*(1), 12–16. <https://doi.org/10.13057/biofar/f090103>
- Patonah, P., Susilawati, E., & Riduan, A. (2018). Aktivitas Antiobesitas Ekstrak Daun Katuk (*Sauropus androgynus* L.Merr) Pada Model Mencit Obesitas. *PHARMACY: Jurnal Farmasi Indonesia (Pharmaceutical Journal of Indonesia)*, *14*(2), 137. <https://doi.org/10.30595/pharmacy.v14i2.1715>
- Pramushinta, I. A. K., Nurhayati, U., & Sukarjati. (2019). *Potensi Ekstrak Etanol Daun Sambung Nyawa (Gynura procumbens), Biji Mahoni (Swietenia mahagoni jacq) Serta Kombinasi Kedua Ekstrak Sebagai Herbal Anti Diabetik Dengan Hewan Coba Mencit (Mus musculus L.)*. 7. <http://snhrp.unipasby.ac.id/>
- Prasetyani, D. & Sodikin. (2017). Analisis Faktor Yang Mempengaruhi Kejadian Diabetes Melitus (DM) Tipe 2. *Jurnal Kesehatan Al Irsyad*, *X*(2), 9.
- Santoso, S. D., & Suryanto, I. (2017). Komparasi Efek Pemberian Minyak Jintan Hitam (*nigella Sativa*) Dengan Minyak Zaitun (*Olea europea*) Terhadap Penurunan Glukosa Darah Pada Mencit (*Mus musculus*) Strain Balb/C. *Jurnal SainHealth*, *1*(1), 36. <https://doi.org/10.51804/jsh.v1i1.76.36-42>
- Scheithauer, T. P. M., Dallinga-Thie, G. M., de Vos, W. M., Nieuwdorp, M., & van Raalte, D. H. (2016). Causality of Small and Large Intestinal Microbiota in Weight Regulation and Insulin Resistance. *Molecular Metabolism*, *5*(9), 759–770. <https://doi.org/10.1016/j.molmet.2016.06.002>
- Serván, P. R. (2013). Obesity and Diabetes. *Nutricion Hospitalaria*, *28*(5), 138–143.
- Sinata, N., & Arifin, H. (2016). Antidiabetes dari Fraksi Air Daun Karamunting (*Rhodomyrtus tomentosa* (Ait.) Hassk.) Terhadap Kadar Glukosa Darah Mencit Diabetes. *Jurnal Sains Farmasi & Klinis*, *3*(1), 72. <https://doi.org/10.29208/jsfk.2016.3.1.102>
- Suwanto, S., & Rahmawati, R. (2019). Aktivitas Hipoglikemik Diet Pakan Ekstrak Biji Labu Kuning (*Cucurbita moschata* Duch) Pada Mencit Diabetes Melitus Terpapar Streptozotocin. *JPSCR: Journal of Pharmaceutical Science and Clinical Research*, *4*(1), 39. <https://doi.org/10.20961/jpscr.v4i1.27292>
- Tillman, E. J., Morgan, D. A., Rahmouni, K., & Swoap, S. J. (2014). Three Months of High-Fructose Feeding Fails to Induce Excessive Weight Gain or Leptin Resistance in Mice. *PLoS ONE*, *9*(9), e107206. <https://doi.org/10.1371/journal.pone.0107206>
- Wilding, J. P. H. (2014). The importance of weight management in type 2 diabetes mellitus. *International Journal of Clinical Practice*, *68*(6), 682–691. <https://doi.org/10.1111/ijcp.12384>

**THIS PAGE INTENTIONALLY LEFT BLANK**